

Self-lensing of a Singular Isothermal Sphere

Yun Wang

*Dept. of Physics, 225 Nieuwland Science Hall, University of Notre Dame, Notre Dame, IN
46556-5670*

(February 18, 2000)

Abstract

Many astrophysical systems can be approximated as isothermal spheres. In an isothermal sphere, the “foreground” objects can act as lenses on “background” objects in the same distribution. We study gravitational lensing by a singular isothermal sphere analytically. Our results may have interesting applications.

PACS numbers: 98.62.Sb, 95.30.Sf

I. INTRODUCTION

Many astrophysical systems can be approximately described as isothermal spheres, e.g. globular clusters, elliptical galaxies, clusters of galaxies. They usually consist (at least partly) of smaller “particles” which are more compact objects, like stars or galaxies. Seen from the outside, i.e. in projection, light from the “background” particles has to pass near the “foreground” particles, and hence it is affected by their gravitational attraction. This effect of “self-lensing” by a stellar disk has been studied in the past [1]. The self-lensing by an isothermal sphere is a simple, well defined problem which may have interesting applications. In this paper, we study the gravitational lensing of the constituents of a singular isothermal sphere (SIS) analytically. For simplicity of notation, we refer to the objects which constitute the SIS as “stars”.

In §2 we define some expressions which are used in the later sections. In §3 we give analytical expressions of the optical depths for self-lensing by a SIS, the number of lensed stars, and the duration distribution of lensing events. In §4, we derive the expressions for the number of lensed stars for self-lensing by a SIS, assuming an arbitrary distribution in the luminosity of stars. The special case of a power law distribution in the luminosity of stars is treated in §5. In §6, we apply our results to the centers of the Milky Way and M87. We give a brief summary in §7. Appendixes A and B contain results for a SIS lensing background sources (located beyond the cut-off radius of the SIS) for comparison.

II. BASIC EQUATIONS

Here we give the basic equations and definitions which are used in later sections. A singular isothermal sphere (SIS) has the mass density

$$\rho_L(r) = \rho_L(0) \left(\frac{r_c}{r} \right)^2 = \frac{M_g}{4\pi r_g} \cdot \frac{1}{r^2} \quad (1)$$

at distance r from the center, where $\rho_L(0)$ and r_c are constants, and M_g is the mass enclosed within radius r_g . Let us assume that the SIS consists of only stars with mass M_L , with the number density $n_L(r) = \rho_L(r)/M_L$.

We consider a SIS with a cutoff radius r_g and its center located at distance r_0 away from the observer. The angular Einstein radius θ_E of an individual star is given by [2]

$$\theta_E^2 = \frac{4GM_L}{c^2} \cdot \frac{D_{ds}}{D_d D_s}, \quad (2)$$

where D_s and D_d are the distances of observer to source and lens, respectively, and D_{ds} is the distance between the lens and the source. Fig. 1 illustrates the geometry between observer, lenses and sources in the SIS. The following expressions will be useful:

$$\begin{aligned} r_d &= \sqrt{r_0^2 + D_d^2 - 2r_0 D_d \cos \theta}, \\ r_s &= \sqrt{r_0^2 + D_s^2 - 2r_0 D_s \cos \theta}, \\ r_1 &= r_0 \cos \theta - \sqrt{r_g^2 - r_0^2 \sin^2 \theta}, \end{aligned}$$

$$\begin{aligned}
r_2 &= r_0 \cos \theta + \sqrt{r_g^2 - r_0^2 \sin^2 \theta}, \\
\alpha &\equiv \frac{r_g}{r_0}, \\
\alpha_0 &\equiv \sqrt{\alpha^2 - \sin^2 \theta}
\end{aligned} \tag{3}$$

where θ is the angle between the line of sight to the source and the line of sight to the center of the SIS; r_d and r_s are the lens' and the source's distances from the center of the SIS respectively; r_1 and r_2 are respectively the smallest and largest possible distances from the observer to a star in the SIS for given θ .

The optical depth for gravitational lensing for a source located in the direction θ and at a distance of D_s from the observer is¹

$$\tau(D_s, \theta) = \int_{D_{d,min}}^{D_{d,max}} dD_d \pi [D_d \theta_E]^2 n_L(r_d), \tag{4}$$

where $D_{d,max}$ and $D_{d,min}$ are the maximum and minimum distances to a lens respectively. The angular number density of lensed sources is

$$\frac{dN_L(\theta)}{d\theta} = 2\pi \sin \theta \int_{D_{s,min}}^{D_{s,max}} dD_s n_s(r_s) D_s^2 \tau(D_s, \theta) \tag{5}$$

where $D_{s,max}$ and $D_{s,min}$ are the maximum and minimum distances to a source respectively. This should be compared with the angular number density of sources

$$\frac{dN(\theta)}{d\theta} = 2\pi \sin \theta \int_{D_{s,min}}^{D_{s,max}} dD_s n_s(r_s) D_s^2. \tag{6}$$

For a given direction θ , the probability of finding a source which is lensed is

$$P(\theta) = \frac{dN_L(\theta)/d\theta}{dN(\theta)/d\theta}. \tag{7}$$

Again, “lensed” here means the source lies inside the Einstein radius of a foreground object.

III. ANALYTICAL RESULTS

In this section, we calculate the optical depth for gravitational self-lensing by a singular isothermal sphere (SIS), the number of lensed stars, and the duration distribution of lensing events. Both lenses and sources are distributed throughout the SIS (each constituent of the singular isothermal sphere can act both as a lens and as a source). The optical depth for gravitational lensing for a source located at a distance of D_s from the observer is

¹Here we employ the definition for “optical depth” in the sense that it is the fractional area that is covered by the Einstein circles of the lenses, that means a minimum magnification of $\mu = 1.34$ is required, see Paczynski (1986) [3].

$$\begin{aligned}
\tau(D_s, \theta) &= \int_{r_1}^{D_s} dD_d \pi [D_d \theta_E]^2 n_L(r_d) \\
&= \frac{GM_g}{c^2 r_g} \left\{ \frac{r_1}{D_s} - 1 + \frac{1}{2} \left(1 - \frac{2r_0 \cos \theta}{D_s} \right) \ln \left[\frac{(D_s - r_0 \cos \theta)^2 + r_0^2 \sin^2 \theta}{r_g^2} \right] \right. \\
&\quad \left. + \frac{D_s \cos \theta - r_0 \cos 2\theta}{D_s \sin \theta} \left[\arctan \left(\frac{D_s - r_0 \cos \theta}{r_0 \sin \theta} \right) + \arctan \left(\frac{\alpha_0}{\sin \theta} \right) \right] \right\}.
\end{aligned} \tag{8}$$

For $\theta \ll 1$, $\tau(D_s, \theta)$ is dominated by the last term in Eq.(8); we find

$$\begin{aligned}
\tau(D_s > r_0, \theta \ll 1) &\simeq \frac{\pi GM_g}{c^2 r_g} \frac{D_s - r_0}{D_s} \frac{1}{\theta}, \\
\tau(D_s < r_0, \theta \ll 1) &\simeq \frac{GM_g}{c^2 r_g} \left\{ \frac{1}{2} \left(\frac{2r_0}{D_s} - 1 \right) \ln \left[\frac{r_g^2}{(D_s - r_0)^2 + D_s r_0 \theta^2} \right] + \frac{1}{D_s} \left[\frac{r_0 - D_s}{\alpha} - r_g \right] - 1 \right\}.
\end{aligned} \tag{9}$$

Note that because the lens number density $n(r_d) \propto 1/r_d^2$, for $D_s < r_0$ (source is closer to the observer than the center of the SIS), the lens number density is finite everywhere, $\tau(D_s, \theta)$ is roughly independent of θ ; while for $D_s > r_0$, the lens number density is infinite at $D_d = r_0$ and $\theta = 0$, $\tau(D_s, \theta)$ is roughly proportional to $1/\theta$. In Fig.2, we show the gravitational lensing optical depth in the galactic center region due to self-lensing, assuming that the stellar cluster there is described by a SIS. Fig.2 shows $\bar{\tau}(D_s, \theta) = \tau(D_s, \theta) [GM_g/(c^2 r_g)]^{-1}$ for $r_0 = 8.5$ kpc, as function of D_s for $\theta = 0.1''$, $1''$, and $10''$, for SIS cutoff radius $r_g = 100$ pc (solid lines) and 1 kpc (dotted lines) respectively.

We assume that the distribution of sources in the SIS are also of the same form as that of the lenses, i.e., $n_s(r_s) \propto 1/r_s^2$. After integrating Eqs.(5) with the $\tau(D_s, \theta)$ from Eq.(8), and taking $D_{s,min} = r_1$, $D_{s,max} = r_2$ (see Fig.1), we find

$$\begin{aligned}
\frac{dN_L(\theta)}{d\theta} &= 8\pi n_s(0) r_c^2 r_0 \left(\frac{GM_g}{c^2 r_g} \right) \left\{ \sin^2 \theta \arctan \left(\frac{\alpha_0}{\sin \theta} \right) - \alpha_0 \sin \theta + \int_0^{\alpha_0/\sin \theta} dx \frac{x \arctan x}{x^2 + 1} \right\}, \\
\frac{dN(\theta)}{d\theta} &= 4\pi n_s(0) r_c^2 r_0 \left[\alpha_0 \sin \theta + \cos 2\theta \arctan \left(\frac{\alpha_0}{\sin \theta} \right) \right].
\end{aligned} \tag{10}$$

For $\theta \ll 1$, the angular densities of lensed sources N_L and of all sources N become

$$\begin{aligned}
\frac{dN_L(\theta)}{d\theta} &\simeq 8\pi n_s(0) r_c^2 r_0 \left(\frac{GM_g}{c^2 r_g} \right) \left\{ \sin^2 \theta \arctan \left(\frac{\alpha_0}{\sin \theta} \right) - \alpha_0 \sin \theta \right. \\
&\quad \left. + \ln \left(\frac{\alpha}{\sin \theta} \right) \arctan \left(\frac{\alpha_0}{\sin \theta} \right) - \frac{\pi}{2} \ln 2 \right\} \\
&\simeq 4\pi^2 n_s(0) r_c^2 r_0 \left(\frac{GM_g}{c^2 r_g} \right) \ln \left(\frac{\alpha}{2\theta} \right). \\
\frac{dN(\theta)}{d\theta} &= 2\pi^2 n_s(0) r_c^2 r_0
\end{aligned} \tag{11}$$

The number densities of both sources and lensed sources increase rapidly as we go to small angles θ . The total number of lensed sources and of sources within angle θ_0 of the line of sight of the SIS center are

$$\begin{aligned}
N_L(\theta_0) &= 16\pi^3 \frac{G\rho_L(0)}{c^2} n_s(0) r_c^4 r_0 \theta_0 \left[\ln \left(\frac{\alpha}{2\theta_0} \right) + 1 \right], \\
N(\theta_0) &= 2\pi^2 n_s(0) r_c^2 r_0 \theta_0.
\end{aligned} \tag{12}$$

The probability of finding a lensed source in the direction θ is

$$P(\theta) \simeq \frac{2GM_g}{c^2 r_g} \ln \left(\frac{\alpha}{2\theta} \right), \quad \theta \ll 1. \tag{13}$$

Note that $N_L(\theta)$ and $P(\theta)$ are not sensitive to $\alpha = r_g/r_0$, the ratio of the SIS cutoff radius to the observer's distance to the SIS center. Eqs.(11)-(13) are very good approximations for $\alpha \geq 10^{-2}$ and $\theta \leq 20''$.

The duration of a lensing event is given by

$$\Delta t \sim \frac{r_0 \theta_E}{v_d} \propto \sqrt{D_{ds}}, \tag{14}$$

where v_d denotes the relative velocity of the lens with respect to the source. In general, the lenses (and sources) have a distribution of velocities.

From the observational point of view, we would like to know the number of lensing events which are shorter than a given time threshold, say, the time duration of the observation. Only the lensing events with time durations shorter than this threshold can be observed with high certainty. In our context, the number of lensing events with time duration shorter than a given threshold, t_c , translates to the number of lensing events with lens-source separation D_{ds} less than a length threshold $r_t(t_c, v_d)$, with

$$r_t(t_c, v_d) = \frac{c^2}{4GM_L} t_c^2 v_d^2 \equiv A t_c^2 v_d^2. \tag{15}$$

Within angle θ_0 of the line of sight to the SIS center, the number of lensing events with duration shorter than t_c is

$$N_L(\Delta t < t_c, \theta_0) = \int_{-\infty}^{\infty} dv_d p(v_d) N_L(D_{ds} \leq r_t(t_c, v_d), \theta_0), \tag{16}$$

where $p(v_d)$ is the probability distribution function of the lens velocity v_d .

Within angle θ_0 of the line of sight to the SIS center, the number of lensing events with lens-source separation D_{ds} less than r_t is given by

$$\begin{aligned}
N_L(D_{ds} \leq r_t, \theta_0) &= \int_0^{\theta_0} 2\pi \sin \theta d\theta \left\{ \int_{r_1}^{r_2} dD_s \int_{D_s - r_t}^{D_s} dD_d - \int_{r_1}^{r_1 + r_t} dD_s \int_{D_s - r_t}^{r_1} dD_d \right\} \\
&\quad D_s^2 n_s(r_s) \pi (D_d \theta_E)^2 n_L(r_d)
\end{aligned} \tag{17}$$

If $r_0 \theta_0 \ll r_t \ll r_g$, we find

$$N_L(D_{ds} \leq r_t, \theta_0) \simeq 16\pi^3 \frac{G\rho_L(0)}{c^2} n_s(0) r_c^4 r_0 \theta_0 \left[\ln \left(\frac{r_t}{2r_0 \theta_0} \right) + 1 \right]. \tag{18}$$

For any realistic distributions of the lens velocities v_d , say, a Gaussian distribution, Eq.(16) has to be solved numerically. To obtain an order of magnitude estimate, we set the

lens velocity to $v_d \sim v_0 = \sqrt{4\pi G \rho_L(0) r_c^2}$. Hence, the number of lensing events with time duration shorter than t_c , within angle θ_0 is

$$N_L(\Delta t < t_c, \theta_0) \sim N_L(D_{ds} \leq r_t(t_c, v_0), \theta_0) \\ \simeq 16\pi^3 \frac{G \rho_L(0)}{c^2} n_s(0) r_c^4 r_0 \theta_0 \left[\ln \left(\frac{A t_c^2 v_0^2}{2 r_0 \theta_0} \right) + 1 \right], \quad (19)$$

where the second approximation follows from Eq.(18), and is only valid for $r_0 \theta_0 \ll r_t(t_c, v_0) = A t_c^2 v_0^2 \ll r_g$, i.e., for lensing events with durations which are neither too long, nor too short. Comparing Eq.(18) with Eq.(12), we find that 50% of all lensing events have $D_{ds} \leq r_t$, with $r_0 \theta_0 \ll r_t = 0.86 \sqrt{r_g r_0 \theta_0} \ll r_g$. This indicates that Eq.(19) is a useful approximation.

Note that we have assumed that all lenses have the same mass for simplicity. The distribution in the lens mass, as well as the distribution in lens/source velocities will contribute to the broadening of the Δt distribution.

IV. NUMBER OF LENSED SOURCES FOR AN ARBITRARY LUMINOSITY FUNCTION

Observations are usually flux-limited, sources with apparent flux below the flux limit of the observation will not be detected. All known astrophysical sources have a distribution in their intrinsic luminosities. The apparent flux of a source is determined by three factors: the intrinsic luminosity of the source, the distance of the source, and the magnification of the source due to gravitational lensing. In this section, we examine the effect of all three factors on the number of lensed sources, allowing for an arbitrary distribution in the intrinsic luminosity of the sources, since luminosity functions are generally poorly known.

Let $\phi(M)$ be the luminosity function of the stars. The fraction of stars with intrinsic luminosity greater than f_0 is

$$Q_0(f_0) = \frac{\int_{-\infty}^{M_0} dM \phi(M)}{\int_{-\infty}^{\infty} dM \phi(M)}, \quad (20)$$

M_0 is the absolute magnitude corresponding to f_0 . $Q_0(f_1) = 1$ defines the low end of the luminosity distribution, f_1 . In general, the inclusion of a distribution in the luminosity of stars leads to a reduction in the expected number of lensed stars.

We write the apparent luminosity of a star with and without gravitational lensing as

$$f_L = \frac{C_1 \mu f_0}{D_s^2}; \quad f_{NL} = \frac{C_1 f_0}{D_s^2}, \quad (21)$$

where C_1 is a constant, μ is the magnification, f_0 is the intrinsic luminosity of the star, and D_s is the observer's distance to the star. The fractions of stars with apparent luminosity greater than f_{min} with and without gravitational lensing are

$$Q_L(f_{min}) = Q_0 \left(\frac{D_s^2 f_{min}}{C_1 \mu} \right), \quad Q_{NL}(f_{min}) = Q_0 \left(\frac{D_s^2 f_{min}}{C_1} \right) \quad (22)$$

The number of sources with apparent luminosity greater than f_{min} without gravitational lensing within angle θ_0 from the line of sight to the SIS center is

$$N(\theta_0, f_{min}) = 2\pi \int_0^{\theta_0} d\theta \sin \theta \int_{D_{s,min}}^{D_{s,max}} dD_s D_s^2 n_s(r_s) Q_0(D_s^2 f_{min}/C_1). \quad (23)$$

For gravitational lensing by the SIS, the number of lensed sources with apparent luminosity greater than f_{min} within angle θ_0 is

$$N_L(\theta_0, f_{min}) = 2\pi \int_0^{\theta_0} d\theta \sin \theta \int_{D_{s,min}}^{D_{s,max}} dD_s D_s^2 n_s(r_s) \int_{f_1}^{\infty} df_0 \left| \frac{dQ_0(f_0)}{df_0} \right| \sigma(D_s, f_0, \mu), \quad (24)$$

where $\sigma(D_s, f_0, \mu)$ is the probability of lensing with magnification greater than $\mu = f_{min}/f_{NL} = f_{min}D_s^2/(f_0C_1)$,

$$\sigma(D_s, f_0, \mu) = \int_{D_{d,min}}^{D_s} dD_d \pi [D_d \theta_E \chi(\mu)]^2 n_L(r_d) = \tau(D_s, \theta) \chi^2(\mu). \quad (25)$$

where $\chi(\mu)$ is the dimensionless angular position of the source relative to the optical axis (line connecting the observer and lens) in units of θ_E for magnification μ . Magnifications greater than μ require $\chi < \chi(\mu)$.

Gravitational lensing by a point mass lens leads to two images with magnifications

$$\mu_{\pm} = \frac{1}{2} \left(1 \pm \frac{\chi^2 + 2}{\chi \sqrt{\chi^2 + 4}} \right). \quad (26)$$

When the lens mass is of order $M_{\odot}$ (and the distances involved are large), the resultant images have micro arcsec scale separation and can not be resolved. For unresolved images, we have

$$\mu = \mu_+ + \mu_- = \frac{\chi^2 + 2}{\chi \sqrt{\chi^2 + 4}}. \quad (27)$$

Hence we write

$$\chi^2(\mu) = \begin{cases} 2 \left(\frac{\mu}{\sqrt{\mu^2 - 1}} - 1 \right), & \mu > \mu_c, \\ 1, & \mu \leq \mu_c, \end{cases} \quad (28)$$

where $\mu_c = 3/\sqrt{5}$. This formula underestimates $\chi^2(\mu)$ for μ close to 1.

For resolved images, a lensed source can be detected if its brighter image (with magnification μ_+) has apparent luminosity greater than f_{min} . $\chi^2(\mu_+)$ is given by Eq.(28), with μ replaced by $2\mu_+ - 1$ on the right hand side of the equation and $\mu_c = (3/\sqrt{5} + 1)/2$. This would apply to the lensing of stars by a central massive black hole; we mention it here for completeness.

Using Eq.(28), we find

$$N_L(\theta_0, f_{min}) = 2\pi \int_0^{\theta_0} d\theta \sin \theta \int_{D_{s,min}}^{D_{s,max}} dD_s D_s^2 n_s(r_s) \tau(D_s, \theta) Q(f_{min}, D_s), \quad (29)$$

with

$$\begin{aligned} Q(f_{min}, D_s) &= \int_{f_1}^{\infty} df_0 \left| \frac{dQ_0(f_0)}{df_0} \right| \chi^2(\mu) \\ &= 2 \int_{x_1}^{1/\mu_c} dx \frac{x}{(1-x^2)^{3/2}} Q_0(x) + 2 \left(\frac{1}{\sqrt{1-x_1^2}} - 1 \right), \end{aligned} \quad (30)$$

where

$$x = \frac{f_0 C_1}{f_{min} D_s^2}, \quad x_1 = \frac{f_1 C_1}{f_{min} D_s^2}. \quad (31)$$

Note that $Q_0(x) = Q_0(f_0)$, and f_1 is defined by $Q_0(f_1) = 1$.

The total number of lensed sources with unlensed apparent luminosity greater than f_{min} (i.e., they can be detected without the help of lensing) is

$$N_L(\theta_0, f_{min}, f_{NL} > f_{min}) = 2\pi \int_0^{\theta_0} d\theta \sin \theta \int_{D_{s,min}}^{D_{s,max}} dD_s D_s^2 n_*(r_s) \tau(D_s, \theta) Q_0(D_s^2 f_{min}/C_1). \quad (32)$$

This gives us the number of sources which are detected due to gravitational lensing

$$N_L(\theta_0, f_{min}, f_{NL} < f_{min}) = N_L(\theta_0, f_{min}) - N_L(\theta_0, f_{min}, f_{NL} > f_{min}). \quad (33)$$

When the dimension of the source distribution is much smaller than our distance to the SIS center (i.e., $r_g \ll r_0$ for self-lensing of SIS), we can take $C_1 = \overline{D_s}^2$, where $\overline{D_s}$ is the average distance to the sources ($\overline{D_s} = r_0$ in the case of SIS self-lensing), then $x_1 \simeq f_1/f_{min}$, $Q_0(D_s^2 f_{min}/C_1) \simeq Q_0(f_{min})$, and $Q(f_{min}, D_s) \simeq Q(f_{min})$. The distribution in the intrinsic luminosities of the sources modifies the total number of observed stars, and the total number of observed stars lensed by other stars (within angle θ_0 of the line of sight to the center of the SIS, and with apparent flux greater than f_{min}) as follows

$$\begin{aligned} N(\theta_0, f_{min}) &\simeq Q_0(f_{min}) N(\theta_0), \\ N_L(\theta_0, f_{min}) &\simeq Q(f_{min}) N_L(\theta_0), \\ N_L(\theta_0, f_{min}, f_{NL} > f_{min}) &\simeq Q_0(f_{min}) N_L(\theta_0), \\ N_L(\theta_0, f_{min}, f_{NL} < f_{min}) &\simeq [Q(f_{min}) - Q_0(f_{min})] N_L(\theta_0), \end{aligned} \quad (34)$$

where $N(\theta_0)$ and $N_L(\theta_0)$ are given in the previous section.

V. NUMBER OF LENSED SOURCES FOR A POWER-LAW LUMINOSITY FUNCTION

The simplest form for the fraction of stars with intrinsic luminosity greater than f_0 is

$$Q_0(f_0) = \begin{cases} \left(\frac{f_0}{f_1} \right)^{-\beta}, & \text{for } f_0 > f_1; \\ 1, & \text{for } f_0 < f_1, \end{cases} \quad (35)$$

where β is a positive constant.

The fractions of stars with apparent luminosity greater than f with and without gravitational lensing are

$$\begin{aligned} Q_L(f) &= \begin{cases} \left(\frac{D_s^2 f}{C_1 \mu f_1}\right)^{-\beta}, & \text{if } \frac{D_s^2 f}{C_1 \mu} > f_1; \\ 1, & \text{else.} \end{cases} \\ Q_{NL}(f) &= \begin{cases} \left(\frac{D_s^2 f}{C_1 f_1}\right)^{-\beta}, & \text{if } \frac{D_s^2 f}{C_1} > f_1; \\ 1, & \text{else.} \end{cases} \end{aligned} \quad (36)$$

The number of sources with apparent luminosity greater than f_{min} without gravitational lensing within angle θ_0 of $\theta = 0$ is

$$N(\theta_0, f_{min}) = 2\pi \left(\frac{f_1}{f_{min}}\right)^\beta \int_0^{\theta_0} d\theta \sin \theta \int_{D_{s,min}}^{D_{s,max}} dD_s D_s^2 \left(\frac{C_1}{D_s^2}\right)^\beta n_s(r_s), \quad (37)$$

where we have assumed $f_{min} D_s^2 / C_1 > f_1$.

For gravitational lensing by the SIS, the number of lensed sources with apparent luminosity greater than f_{min} within angle θ_0 of $\theta = 0$ is

$$\begin{aligned} N_L(\theta_0, f_{min}) &= 2\pi \left(\frac{f_1}{f_{min}}\right)^\beta \int_0^{\theta_0} d\theta \sin \theta \int_{D_{s,min}}^{D_{s,max}} dD_s D_s^2 \left(\frac{C_1}{D_s^2}\right)^\beta n_s(r_s) \\ &\quad \cdot \tau(D_s, \theta) F(\beta, f_{min}, D_s), \end{aligned} \quad (38)$$

with

$$F(\beta, f_{min}, D_s) = 2\beta \int_{x_1}^{1/\mu_c} dx \frac{1}{x^{\beta+1}} \left(\frac{1}{\sqrt{1-x^2}} - 1 \right) + \mu_c^\beta, \quad (39)$$

where $x_1 = f_1 C_1 / (f_{min} D_s^2)$. The number of lensed stars with unlensed flux greater than f_{min} is

$$N_L(\theta_0, f_{min}, f_{NL} > f_{min}) = 2\pi \left(\frac{f_1}{f_{min}}\right)^\beta \int_0^{\theta_0} d\theta \sin \theta \int_{D_{s,min}}^{D_{s,max}} dD_s D_s^2 \left(\frac{C_1}{D_s^2}\right)^\beta n_s(r_s) \tau(D_s, \theta). \quad (40)$$

For $x_1 = f_1 C_1 / (f_{min} D_s^2) \ll 1$, we find

$$F(\beta, f_{min}, D_s) = \begin{cases} 2.236 - x_1, & \beta = 1; \\ 1.677 - 2 \ln x_1, & \beta = 2. \end{cases} \quad (41)$$

Note that for power-law $Q_0(f_0)$, $Q_0(f_{min}) = (f_1/f_{min})^\beta$, and $Q(f_{min}) = (f_1/f_{min})^\beta F(\beta, f_{min}, D_s)$. Recall that $Q_0(f_{min})$ is the reduction factor in the total number of observed stars (also the reduction factor in the total number of lensed stars which can be observed without the help of gravitational lensing), $Q(f_{min})$ is the reduction factor in the total number of observed stars lensed by other stars. $[Q(f_{min}) - Q_0(f_{min})] N_L(\theta_0)$ gives the number of stars observed due to gravitational lensing by the SIS.

VI. APPLICATIONS TO THE CENTERS OF THE MILKY WAY AND M87

An obvious application of our results is the self-lensing of the star cluster near the Galactic center. Although the gravitational lensing near the Galactic center is dominated by the central black hole [4], the number of lensed stars due to self-lensing of the star cluster near the Galactic center is only about an order of magnitude smaller than the number of stars lensed by the central black hole. Gravitational lensing studies near the Galactic center can provide independent evidence for the existence of the massive black hole at the Galactic center, and constrain the stellar density near the Galactic center.

We take the distance to the Galactic center to be $r_0 = 8.5$ kpc. For the density profile of the SIS (see Eq.(1)), we take $\rho_L(0) = 4 \times 10^6 M_\odot \text{pc}^{-3}$, and $r_c = 0.17$ pc. In a cone with angular radius θ_0 centered around our line of sight to the Galactic center, without considering the distribution in the luminosity of the stars, the number of lensed stars due to self-lensing of the star-cluster is [see Eq.(12)]

$$N_L^{GC}(\theta_0) = 160\pi^3 \frac{G\rho_L^2(0)r_c^4}{M_L c^2} r_0 \theta_0 \gamma(\theta_0, r_g) \quad (42)$$

$$= 0.1345 \gamma(\theta_0, r_g) \left(\frac{\theta_0}{1''} \right) \left(\frac{M_\odot}{M_L} \right), \quad (43)$$

where r_g is the cutoff radius of the SIS, and $\gamma(\theta_0, r_g) \equiv \ln[r_g/(2r_0\theta_0) + 1]/10$. Taking $\theta_0 = 1''$, for $r_g = 100$ pc, 1 kpc, and 10 kpc, $\gamma = 0.81, 1.04$, and 1.27 respectively.

When we include the distribution in the luminosity of the stars by using the empirical luminosity function from Mamon & Soneira (1982) [5], the number of lensed stars is reduced by approximately one order of magnitude from that given by Eq.(42), for limiting K magnitude of 21 and an extinction of 3 magnitudes to the Galactic center.

The duration of a lensing event is given by

$$\Delta t \sim \frac{r_0 \theta_E}{v_d} \sim 2 \text{ days} \left(\frac{M_L}{M_\odot} \right)^{1/2} \left(\frac{D_{ds}}{\text{pc}} \right)^{1/2} \quad (44)$$

where $v_d = \sqrt{4\pi G \rho_L(0) r_c^2} \simeq 80$ km/s is the mean velocity of the lens, θ_E is the angular Einstein radius given by Eq.(2), and D_{ds} is the separation between the source and the lens.

The number of lensing events with lens-source separation D_{ds} less than r_t within angle θ_0 of the line of sight to the SIS center, $N_L(D_{ds} \leq r_t, \theta_0)$, is given by Eq.(18). 50% of all lensing events have $D_{ds} \leq 0.86 \sqrt{r_g r_0 \theta_0} \ll r_g$ (§3), i.e., roughly 50% of all lensing events have duration shorter than

$$\Delta t \sim 2.64 \text{ days} \left(\frac{\theta_0}{1''} \right)^{1/4} \left(\frac{r_g}{100 \text{ pc}} \right)^{1/4} \left(\frac{M_L}{M_\odot} \right)^{1/2}. \quad (45)$$

The observation of the self lensing of stars near the Galactic center would require substantial observational resources, and should only be a by-product of ultra high resolution proper motion studies of stars near the Galactic center. Another by-product will be observing the lensing of the same stars due to the black hole at the Galactic center. Note that the lensing events due to the central black hole last about three order of magnitude longer than

the self-lensing events. For a given observation with finite duration and dense time sampling (say, the detailed and accurate monitoring of stellar motions near the Galactic center for two years), the smaller number of self-lensing events is balanced by their shorter time durations in favor of their detection.

Note that the number of lensed events increases with both typical lens velocity $v_d = \sqrt{4\pi G\rho_L(0)r_c^2}$, and our distance to the source r_0 [see Eqs.(12) and (42)]. For galaxies which are more distant from us or/and have higher central mass density, the number of lensed stars increases. Our distance to M87 is about 15 Mpc. Within 1 arcsecond (about 73 pc) of the center of M87, the velocity dispersions of stars are about 400 km/s [6]. This implies that M87 has a central stellar density at least as large as that of the Milky Way. For M87, $\gamma(\theta_0, r_g) = \ln[r_g/(2r_0\theta_0) + 1]/10 \sim 0.1$. Therefore

$$N_L^{M87}(\theta_0) \gtrsim 176.5 N_L^{GC}(\theta_0) \left(\frac{\theta_0}{1''}\right) \left(\frac{M_\odot}{M_L}\right) \sim 24 \left(\frac{\theta_0}{1''}\right) \left(\frac{M_\odot}{M_L}\right). \quad (46)$$

If a large enough number of self-lensed stars are observed near the center of M87, they can be used to constrain the model for the central mass distribution of M87.

VII. SUMMARY

We have studied the self-lensing of a singular isothermal sphere (SIS) analytically. We have derived simple analytical formulas for the optical depth, the number of lensed stars, and the duration distribution of lensing events. We have also derived the expressions for the number of lensed stars assuming an arbitrary distribution in the luminosity of stars, and we have considered the special case of a power law distribution in the luminosity of stars. Application of our results to the gravitational lensing by the star cluster near the Galactic Center gives potentially observable number of lensing events. High resolution monitoring of the central region of M87 may yield dozens of self-lensed stars which can be used to constrain the model for the central mass distribution of M87.

For comparison, we give analytical expressions of the optical depth for a SIS lensing background sources (located beyond the cut-off radius of the SIS) in Appendix A, and the application of these expressions to the gravitational lensing of distant sources by the star cluster near the Galactic Center in Appendix B.

Since many astrophysical systems can be approximated as isothermal spheres, it is possible that some of them can have significant self-lensing; our results should be useful in studying gravitational lensing in these systems.

ACKNOWLEDGMENTS

It is a pleasure for me to thank Ed Turner for suggesting this calculation; Joachim Wambsganss for providing Figure 1, helpful discussions, and careful readings of the manuscript; and the referee for helpful suggestions.

Appendix A. SIS lensing a background source

For a source located beyond the cutoff radius of the SIS (on the other side of the SIS, away from the observer) and is lensed by the SIS, $D_s > r_2$ (see Fig.1), the optical depth is given by

$$\begin{aligned}\tau(D_s, \theta) &= \int_{r_1}^{r_2} dD_d \pi [D_d \theta_E]^2 n_L(r_d), \\ &= \frac{2GM_g}{c^2 r_g} \left\{ \frac{D_s \cos \theta - r_0 \cos 2\theta}{D_s \sin \theta} \arctan \left(\frac{\alpha_0}{\sin \theta} \right) - \frac{r_0}{D_s} \alpha_0 \right\}.\end{aligned}\quad (47)$$

For a source far behind the SIS, $D_s \gg r_2$, we find

$$\tau(D_s \gg r_2, \theta) = \frac{2GM_g}{c^2 r_g} \cot \theta \arctan \left(\frac{\alpha_0}{\sin \theta} \right), \quad (48)$$

which is independent of D_s ; this is consistent with the trend shown by Fig.2. For $\theta \ll 1$, we find

$$\tau(D_s, \theta \ll 1) \simeq \frac{\pi GM_g}{c^2 r_g} \frac{D_s - r_0}{D_s} \frac{1}{\theta}. \quad (49)$$

Let us assume that the cluster of sources is located behind the SIS, at a mean distance of $\overline{D_s}$ from the center of the SIS, and that the radius of the distribution along the line of sight is r_s . If $\overline{D_s} \gg r_s$, we can take $n_s(D_s) \simeq n_s(\overline{D_s})$. The angular densities of lensed sources and of all sources become

$$\begin{aligned}\frac{dN_L(\theta)}{d\theta} &= 8\pi n_s(\overline{D_s}) \overline{D_s} r_s r_0 \left(\frac{GM_g}{c^2 r_g} \right) \left\{ \left[\frac{\overline{D_s}}{r_0} \cos \theta - \cos 2\theta \right] \arctan \left(\frac{\alpha_0}{\sin \theta} \right) - \alpha_0 \sin \theta \right\}, \\ \frac{dN(\theta)}{d\theta} &= 4\pi n_s(\overline{D_s}) \overline{D_s}^2 r_s \sin \theta.\end{aligned}\quad (50)$$

For $\theta \ll 1$, the angular densities of lensed sources and of all sources become

$$\begin{aligned}\frac{dN_L(\theta)}{d\theta} &\simeq 4\pi^2 n_s(\overline{D_s}) \overline{D_s} (\overline{D_s} - r_0) r_s \left(\frac{GM_g}{c^2 r_g} \right), \\ \frac{dN(\theta)}{d\theta} &= 4\pi n_s(\overline{D_s}) \overline{D_s}^2 r_s \theta.\end{aligned}\quad (51)$$

The probability of finding a lensed source in the direction θ is (see Eq.(7))

$$P(\theta) \simeq \frac{GM_g}{c^2 r_g} \frac{\overline{D_s} - r_0}{\overline{D_s}} \frac{1}{\theta}, \quad \theta \ll 1. \quad (52)$$

Note that although the optical depth in the case of SIS lensing a background source has the same small θ limit as the optical depth in the case of SIS lensing itself, the angular number density of lensed sources increases more rapidly as θ decreases in the case of SIS self-lensing due to the increase in the number of sources.

Appendix B. Lensing of distant sources by the star cluster near the Galactic Center

Again we take SIS density profile for the star cluster near the Galactic center, with $\rho_L(0) = 4 \times 10^6 M_\odot \text{pc}^{-3}$ and $r_c = 0.17 \text{pc}$. For a cluster of sources located on the other side of the Galaxy, we find

$$N_L(\theta_0) = 0.2 \left(\frac{\theta_0}{1''} \right) \left(\frac{n_s(\overline{D_s})}{1 \text{pc}^{-3}} \right) \left(\frac{\overline{D_s}}{17 \text{kpc}} \right) \left(\frac{\overline{D_s} - r_0}{8.5 \text{kpc}} \right) \left(\frac{r_s}{100 \text{pc}} \right). \quad (53)$$

This number should probably decrease by more than an order of magnitude when the luminosity distribution of stars and dust extinction are taken into account.

The duration of a lensing event is given by

$$\Delta t \sim \frac{\overline{D_s} \theta_E}{v_s} \sim 8 \text{ months} \left(\frac{M_L}{M_\odot} \right)^{1/2} \left(\frac{80 \text{km/s}}{v_s} \right) \left(\frac{\overline{D_s}}{17 \text{kpc}} \right)^{1/2} \left(\frac{\overline{D_s} - r_0}{8.5 \text{kpc}} \right)^{1/2} \quad (54)$$

The durations of lensing events are clustered around the typical value because all the sources are at about the same distance behind the lens.

REFERENCES

- [1] Gould, A. 1995, *Astrophys. J.* , 441, 77
- [2] Schneider, P., Ehlers, J., & Falco, E.E. 1992, “Gravitational Lenses”, (Springer-Verlag, Berlin)
- [3] Paczynski, B. 1986, *Astrophys. J.* , 304, 1
- [4] Wardle, M. & Yusef-Zadeh, F. 1992, *Astrophys. J.* , 387, L65
- [5] Mamon, G. & Soneira, R.M. 1982, *ApJ*, 255, 181
- [6] Dehnen, W., in *Proceedings of the Ringberg workshop on M87* (astro-ph/9802224).

FIGURES

FIG. 1. Cartoon illustrating the geometry between observer (O), lenses (L) and sources (S) in the singular isothermal sphere (SIS, represented by the circle centered at C with radius r_g). θ is the angle between the line of sight to the source (OS) and the line of sight to the center of the SIS (OC); the distances $r_d = LC$, $r_s = SC$, $D_d = OL$, $D_{ds} = LS$, $D_s = OS$. r_1 and r_2 are, respectively, the smallest and largest possible distances from the observer (O) to an object in the SIS for given θ .

FIG. 2. Optical depth for self lensing by a singular isothermal sphere (SIS) at the Galactic center (at a distance of 8.5 kpc), $\bar{\tau}(D_s, \theta) = \tau(D_s, \theta) [GM_g/(c^2 r_g)]^{-1}$, as function of the distance to the source D_s , for source direction angle (measured from the line of sight to the Galactic center) $\theta = 0.1''$, $1''$, and $10''$, and for two values of the cut-off radius of the SIS: $r_g = 100$ pc (solid lines) and 1 kpc (dotted lines).



